

Hermitian matrices and cohomology of Kähler varieties

Andrea Causin and Gian Pietro Pirola *

Abstract

We give some upper bounds on the dimension of the kernel of the cup product map $H^1(X, \mathbb{C}) \otimes H^1(X, \mathbb{C}) \rightarrow H^2(X, \mathbb{C})$, where X is a compact Kähler variety without Albanese fibrations.

AMS (MOS) Subject Classification: 32J27 (32J25).

Key words: Kähler varieties, cohomology, fundamental group, hermitian matrices, determinantal varieties.

Introduction

One of the special features of the Kähler geometry is the interplay between topology and linear algebra. The goal of this paper is to give some applications of results in advanced linear algebra appearing in [2] and [14] to obtain bounds on the dimension of the kernel of the cup product mapping

$$\phi: \bigwedge^2 H^1(X, \mathbb{C}) \rightarrow H^2(X, \mathbb{C}),$$

in the case where X is a compact Kähler variety admitting no Albanese fibration.

Recall that the notion of *no Albanese fibration*, for an n -dimensional variety X as before, can be given by requiring that, for any $k < n$ and any independent $\beta_1, \dots, \beta_k \in H^0(\Omega^1(X))$, the product

$$\beta_1 \wedge \dots \wedge \beta_k \in H^0(\Omega^k(X)) = H^{k,0}(X)$$

is not zero. Also remark that $\bigwedge_{i=1}^k \beta_i = 0$ gives rise to an integrable distribution and to a foliation with closed leaves; this allows to define a fibration (see [10] or 1.2.2 later in this paper), hence the definition is consistent.

The main result we prove is:

Theorem 1 *Let X be a compact Kähler variety without Albanese fibrations, and let $\phi: \bigwedge^2 H^1(X, \mathbb{C}) \rightarrow H^2(X, \mathbb{C})$ be the cup product.*

1. *If $q \leq 2n - 1$, then ϕ is injective.*
2. *If $q = 2n$, then $\dim \ker \phi \leq 2c + 3$ where $q = 2^c(2b + 1)$, and b, c integers.*
3. *If $q = 5$ and $n = 2$, then $\dim \ker \phi \leq 14$.*

*Partially supported by 1) PRIN 2005 “Spazi di moduli e teorie di Lie”; 2) Indam (GNSAGA); 3) Far 2006 (PV): “Varietà algebriche, calcolo algebrico, grafi orientati e topologici”.

The previous bounds are achieved thanks to the Hodge decomposition that we shortly recall. We have

$$H^1(X, \mathbb{C}) = H^{1,0} \oplus H^{0,1}; \quad H^2(X, \mathbb{C}) = H^{2,0} \oplus H^{1,1} \oplus H^{0,2},$$

hence ϕ defines maps:

1. $\phi^{2,0} : \bigwedge^2 H^{1,0} \longrightarrow H^{2,0}$ (and its dual $\phi^{0,2}$);
2. $\phi^{1,1} : H^{1,0} \otimes H^{0,1} \longrightarrow H^{1,1}$.

The role of the assumption of no Albanese fibration becomes clear when dealing with the estimate of $\dim \ker \phi^{2,0}$ (remark, for instance, that there are not decomposable elements in the kernel).

We look, then, for some upper-bound for $\kappa = \dim \ker \phi^{1,1}$. Set $H^{1,1}(X)_{\mathbb{R}} = H^{1,1}(X) \cap H^2(X, \mathbb{R})$ and call $M \subset H^{1,0} \otimes H^{0,1}$ the subspace of forms invariant under complex conjugation. The space M is naturally identified with the space $\mathbb{H}_q$ of Hermitian $q \times q$ matrices and $\phi^{1,1}$ restricts to $\phi_{\mathbb{R}}^{1,1} : M \rightarrow H^{1,1}(X)_{\mathbb{R}}$; putting $K = \ker \phi_{\mathbb{R}}^{1,1}$ we have that $\kappa = \dim K$. Assuming that X has no Albanese fibration and using a positivity argument we find restrictions on the signature and hence on the rank of the involved matrices: if $A \in K$, $A \neq 0$, then the rank of A must be $\geq 2n$. In particular when $n = 2q$ then A is invertible and from [2] we achieve a very good bound on κ . Write $q = 2^c(2b+1)$, with b and c integers; we have $\kappa \leq 2c+1$. The basic remark is that the eigenvalues of iA , with $i^2 = -1$, are not real, so if $v \in S^{2q-1} \subset \mathbb{C}^q$ the tangential projection of iAv on the unitary sphere defines a vector field not vanishing at any point. The result is then based on [1].

Unfortunately when $2n < q$ no good bound seems to be known in general. However, at least for symmetric matrices, some very interesting works have been done; among the others, we suggest [13, 14]. In a very strict sense this is a *comeback of algebraic geometry*. The basic idea is to consider the degeneracy locus $\{A : \text{rank } A \leq k\}$ of the matrices as real varieties, then to study its intersection with real subspaces. In particular in [14], after the natural projectivization, an elegant Lefschetz fixed point argument is used on a suitable complex algebraic variety. This shows, in the case of 5×5 real symmetric matrices of rank lower than 4, that the intersection is not empty. By performing the same kind of computations, for $n = 2$ and $q = 5$ we obtain $\kappa \leq 8$.

We would like to focus the importance of the mapping ϕ in the case of compact Kähler varieties. On one side we have Castelnuovo-De Franchis-type theorem concerning Albanese fibrations (see [10]). On the other side, formality theorems [12] imply that the De Rham fundamental group $\pi_1(X) \otimes \mathbb{C}$ is determined by ϕ [4]. We also recall works of Campana (see [9]) on Kähler nilpotent groups.

The paper is organized in the following way. In section 1 we recall the Albanese variety and the Castelnuovo-de Franchis theory as developed by Catanese [10]. In the second section we study the 1,1 real forms and their connection with hermitian matrices. This allows to prove the first two assertions of the above theorem. In section 3 we perform the computations necessary for the first case not covered in [2].

Both authors are delighted to thank Margherita for her helpful contribution and support.

1 Variety of Albanese type

1.1 Albanese variety

Let X be a complex compact Kähler variety of dimension n , and

$$H^{p,q} = H^q(X, \Omega^p)$$

be the Hodge spaces. We have the Hodge decomposition:

$$H^n(X, \mathbb{C}) = \bigoplus_{p+q=n} H^{p,q}.$$

Set $V = H^{1,0}$ and $H^{0,1} = \overline{V}$ its conjugate. Integration defines then

$$j : H_1(X, \mathbb{Z}) \longrightarrow V^*,$$

where $*$ stands for dual. Let

$$Alb(X) = V^*/j(H_1(X, \mathbb{Z}))$$

be the Albanese variety. The irregularity of X is denoted by $q_X = \dim V = \dim A$. The choice of a base point $p \in X$ defines the Albanese map

$$\alpha : X \longrightarrow Alb(X).$$

Definition 1.1.1. *We say that X is of Albanese type if α is generically finite. We say that X is of Albanese strict type if, moreover, α is not surjective. That is:*

$$\dim(\alpha(X)) = \dim(X) < q_X.$$

In the sequel we will assume X of Albanese strict type. For our purposes, thanks to the result of Campana, we could also assume that α is generically one-to-one (see [9] and [4, Ch. 2, Sect. 4]).

1.2 Decomposable form and Albanese fibrations

We describe some Castelnuovo-de Franchis-type theorems. With the previous notations, the cohomology map induced by the Albanese morphism: $\alpha^* : H^k(A, \mathbb{C}) \rightarrow H^k(X, \mathbb{C})$ is a Hodge structure map. We can moreover make the identifications: $H^k(A, \mathbb{C}) \equiv \bigwedge^k H^1(X, \mathbb{C})$, $V = H^{1,0}(A) \equiv H^{1,0}(X)$ and $H^{p,q}(A) = \bigwedge^p V \otimes \bigwedge^q \overline{V}$. We have maps: $\alpha^{p,q} : \bigwedge^p V \otimes \bigwedge^q \overline{V} \rightarrow H^{p,q}$.

Definition 1.2.1. *Given $s > 0$, we say that a (rational map) $f : X \rightarrow Y$ is an s -Albanese fibration if*

1. Y is of Albanese strict type;
2. $\dim X - \dim Y = s$.

When $s = n - 1$, Y is a curve of genus $g > 1$ and f is usually called an irregular pencil.

We have (see [10]):

Proposition 1.2.2. *The following conditions are equivalent*

1. X has no s -Albanese fibration for $s < n - k$
2. $\alpha^{k,0}$ is injective on the decomposable forms:

$$0 \neq \beta_1 \wedge \cdots \wedge \beta_k \in H^{k,0}(X),$$

β_i independent.

Remark 1.2.3. More precisely, Fabrizio Catanese in [10] gave a one-to-one correspondence between fibrations of Albanese type and maximal isotropic subspaces of the first cohomology group of X .

2 Forms and matrices

2.1 Real 1.1 forms

As before, X is of strict Albanese type. We will consider in details the map:

$$\alpha^{1,1} \equiv \phi^{1,1}: H^{1,1}(A) \equiv V \otimes \overline{V} \rightarrow H^{1,1}(X). \quad (1)$$

We set

$$\kappa = \dim \ker(\phi^{1,1}). \quad (2)$$

This is the (1.1) part of $\phi \equiv \alpha^2: H^2(A, \mathbb{C}) \equiv \bigwedge^2 H^1(X, \mathbb{C}) \rightarrow H^2(X, \mathbb{C})$. Since α^2 is a piece of a Hodge structure map, the kernel of α^2 is defined over the rational numbers and a fortiori over the real numbers. We have then a map:

$$\alpha_{\mathbb{R}}^{1,1}: H^{1,1}(A)_{\mathbb{R}} \rightarrow H^{1,1}(X)_{\mathbb{R}} \subset H^2(X, \mathbb{R})$$

and in particular $\kappa = \dim \ker(\alpha_{\mathbb{R}}^{1,1})$. We can identify $H^{1,1}(A)_{\mathbb{R}}$ with the sesquilinear forms on V^* . More explicitly, we fix a basis β_j $j = 1, \dots, q$ of V . An element of $H^{1,1}(A)_{\mathbb{R}}$ has the form $\Omega = i \sum_{j,s} a_{j,s} \beta_j \wedge \overline{\beta_s}$, $i^2 = -1$, $a_{j,s} \in \mathbb{C}$: $\overline{\Omega} = \Omega$, that is the matrix $A(\Omega) = (a_{j,s})$ is hermitian. Let $\mathbb{H}_q$ be the space of $q \times q$ hermitian matrices. We may define $\sigma: \mathbb{H}_q \rightarrow H^{1,1}(X)_{\mathbb{R}}$ as:

$$\sigma(A) = i \sum_{j,s} a_{j,s} \beta_j \wedge \overline{\beta_s}, \quad (3)$$

where $A = (a_{j,s})$. We have $\kappa = \dim \ker \sigma$. In particular, the rank and the signature of a form $\Omega \in H^{1,1}(A)_{\mathbb{R}}$ are well defined.

Definition 2.1.1. If $\Omega \in H^{1,1}(A)_{\mathbb{R}}$ (respectively $A \in \mathbb{H}_q$) has signature (r, s) (and rank $r + s \leq q$), we set $m(\Omega) = \min(r, s)$ (respectively $m(A) = \min(r, s)$).

The following proposition generalizes the elementary result explained in the introduction.

Proposition 2.1.2. Let $k < n$ be an integer, and assume that X has no $n - k$ -Albanese fibration. Let $\Omega \in H^{1,1}(A)_{\mathbb{R}}$, $\Omega \neq 0$. If $m(\Omega) \leq k - 1$ then $\alpha^{1,1}(\Omega) \neq 0$.

Proof. Up to a change between Ω and $-\Omega$, we may assume $s = m(\Omega)$ where (r, s) is the signature of Ω , $s \leq k - 1 < n - 1$. We may find a basis β_i of V such that

$$\Omega = i \sum_{j=1}^r \beta_j \wedge \overline{\beta_j} - i \sum_{j=r+1}^{r+s} \beta_j \wedge \overline{\beta_j} = \Omega^+ - \Omega^-.$$

Setting $\varphi = \beta_{r+1} \wedge \cdots \wedge \beta_{r+s} \in H^{s,0}(X)$ and $\Theta = \varphi \wedge \overline{\varphi}$, we compute:

$$\Omega \wedge \Theta = i \sum_{j=1}^r \beta_j \wedge \overline{\beta_j} \wedge \Theta = (-1)^s i \sum_{j=1}^r \beta_j \wedge \beta_{r+1} \wedge \cdots \wedge \beta_{r+s} \wedge \overline{\beta_j \wedge \beta_{r+1} \wedge \cdots \wedge \beta_{r+s}}.$$

Now, posing $\varphi_j = \beta_j \wedge \beta_{r+1} \wedge \cdots \wedge \beta_{r+s} \in H^{s+1,0}(X)$ and $\Theta_j = \varphi_j \wedge \overline{\varphi_j}$, we have:

$$\Omega \wedge \Theta = i(-1)^s \sum_j \Theta_j.$$

Assume by contradiction $\Omega \in \ker \alpha^{1,1}$. It follows that $\Omega \wedge \Theta = 0$ in $H^{s+1,s+1}(X)$. Fix $\omega = i \sum_{j=1}^q \beta_j \wedge \overline{\beta_j}$; this is the pull-back of a Kähler form on A and is positive on a Zariski open set of X , since the Albanese map of X is generically finite. We get

$$0 = \int_X \Omega \wedge \Theta \wedge \omega^{n-s-1} = \sum_j \int_X i(-1)^s \Theta_j \wedge \omega^{n-s-1}.$$

All terms have the same sign. It follows that

$$\Theta_j \wedge \omega^{n-s-1} = 0;$$

this forces $\Theta_j = 0$ and finally $\varphi_j = \beta_j \wedge \beta_{r+1} \wedge \cdots \wedge \beta_{r+s} = 0$. Since $s+1 \leq k$, we get a contradiction with Prop. 1.2.2. $\square$

We have then the following:

Corollary 2.1.3. *Assume that X has no Albanese fibration and $\Omega \in \ker(\alpha^{1,1})_{\mathbb{R}}$, $\Omega \neq 0$; then, $m(\Omega) \geq n-1$ and $\text{rank}(\Omega) \geq 2n$.*

2.2 Hermitian matrices

Let $\mathbb{H}_q$ be the space of the $q \times q$ hermitian matrices. Let $\mathbb{H}_{q,m} \subset \mathbb{H}_q$ be the subset of the matrices with rank bigger than $m-1$:

$$\mathbb{H}_{q,m} \in \{A \in \mathbb{H}_q : \text{rank}(A) \geq m\}.$$

Definition 2.2.1. *Let $V \subset \mathbb{H}_q$ be a real subspace. We say that V has rank $\geq n$ ($\text{rank}(V) \geq n$) if $V \setminus \{0\} \subset \mathbb{H}_{q,n}$. We set $d_{q,n} = \max_{\text{rank}(V) \geq n} \dim V$.*

Proposition 2.2.2. 1. *If $q \leq m-1$, then $d_{q,m} = 0$.*

2. *$d_{q,q} = 2c+1$, where $q = 2^c(2b+1)$, with b and c integers.*

3. *$d_{5,4} \leq 8$.*

Proof. 1. Obvious.

2. The elements in $\mathbb{H}_{q,q}$ are given by invertible hermitian matrices. Then, we come back to the hermitian case in [2]. This result was obtained as a consequence of [1].

3. To be computed in the next section. $\square$

Recalling that $\kappa = \dim \ker \phi^{1,1} = \dim \ker \sigma$ (see 3), we have the following:

Proposition 2.2.3. *Let $n = \dim X$, $q = \dim H^{1,0}(X)$ and κ as before. Assume that X has no Albanese fibration; then $\kappa \leq d_{q,2n}$.*

Proof. It follows from 2.1.3. □

Here we present some consequences of Prop. 2.2.2:

Corollary 2.2.4. *If $q \leq 2n - 1$, then $\alpha^2: H^2(A, \mathbb{C}) \rightarrow H^2(X, \mathbb{C})$ is injective.*

Proof. Firstly one has that $\alpha^{1,1} \equiv \phi^{1,1}$ is injective. Then consider $\omega \in H^{2,0}(A)$ such that $\alpha^*(\omega) = 0$; we show that $\omega = 0$. Indeed, if it is not, we can find a basis β_i of V for which $\omega = \sum_{i=1}^k \beta_i \wedge \beta_{i+k}$, with $k < n$; taking the $k-1$ form $\phi = \wedge_{i=2}^k \beta_i$, we get $\alpha^*(\omega \wedge \phi) = 0$ and consequently $\wedge_{i=1}^{k+1} \beta_i = 0$ on X . By 1.2.2, this would give an Albanese fibration contradicting our assumptions. □

The previous result is essentially standard linear algebra. The first part of the following proposition is a consequence of the hard topological result of Adams [1, 2].

Proposition 2.2.5. *Assume that X has no Albanese fibration and $q = 2n$. Write $q = 2^c(2b+1)$, b and c being integers. Then we have*

1. $\dim(\ker(\alpha^{1,1})) \leq 2c + 1$ and
2. $\dim(\ker(\alpha^{2,0})) \leq 1$.

The two inequalities above can be unified by saying that $\dim \ker(\alpha^2) \leq 2c + 3$. Consequently it holds: $b_2(X) \geq \dim \operatorname{Im} \phi \geq q(2q-1) - 2c - 3$.

Proof. 1. From 2.2.3 and 2.2.2 we have $\dim \ker \sigma \leq 2c + 1$.

2. Arguing as in 2.1.3, the nontrivial forms in $\ker \alpha^{2,0}$ must be of maximal rank n . Representing them as anti-symmetric matrices, the elements in $\ker \alpha^{2,0} \setminus \{0\}$ are invertible. This is a complex space, and it follows that $\dim(\ker \alpha^{2,0}) \leq 1$. □

Proposition 2.2.6. *Let X be a compact algebraic variety without irregular pencils whose fundamental group admits a presentation with γ generators and ρ relations.*

1. *If $q = 2n$, then $\rho - \gamma \geq q(2q-3) - 2c - 3$; moreover, if X is a surface i.e. $n = 2$, $c_2(X) \geq 7$.*
2. *If $q = 5$ and X is a surface, then $b_2(X) \geq 31$, hence $\rho - \gamma \geq 31$ and $c_2(X) \geq 13$.*

Proof. Firstly recall that (see [3, Th. 1.1] and [4, Ch. 3]) the interplay between the cup product map and the fundamental group give rise to the following estimate: $\rho - \gamma \geq \dim \operatorname{Im}(\phi) - 2q$.

1. The estimate on $\rho - \gamma$ follows directly from the previous remark and 2.2.5. When $q = 2n = 4$, we have $b_1(X) = b_3(X) = 8$ and $b_2(X) \geq 21$.
2. To deduce $b_2(X) \geq 31$ one uses the third part of 2.2.2 together with the fact that $\dim \operatorname{Im} \phi^{2,0} \geq 2q - 3 = 7$. The rest is as in the previous point. □

Remark 2.2.7. *In [3] the following estimate for a compact Kähler variety of any dimension with no irrational pencil is given: $\rho - \gamma \geq 4q - 7$.*

The following result was our historical motivation:

Corollary 2.2.8. *Let X be a minimal projective algebraic surface with $q = 4$ and $p_g = \dim H^{2,0}(X) = 5$. Let K be the canonical bundle of X ; then $16 \leq K^2 \leq 17$.*

Proof. In [6] it was proved that $K^2 \geq 16$ and that if X has an irregular pencil then $K^2 = 16$. When X has no irregular pencil Noether formula (see [7, 15]) $K^2 + c_2(X) = 12\chi_{\text{hol}} = 12(p_g - q + 1) = 24$ and the inequality $c_2(X) \geq 7$ forces then $K^2 \leq 17$. $\square$

Remark 2.2.9. *In the previous example the Miyaoka-Bogomolov-Yau inequality (see for instance [7]) gives only $K^2 \leq 9\chi_{\text{hol}} = 18$. No examples of surfaces with $K^2 = 17$ are actually known.*

3 Real degeneracy loci.

The present section is entirely dedicated to the proof of the third point in 2.2.2. To this purpose, we adapt to hermitian matrices the work developed in [14] in the case of real symmetric matrices, by thinking to the degeneracy locus $\{A \text{ hermitian, } \text{rank} A \leq m\}$ as a real variety. The main theoretic tool proved in [14] is the following theorem, which is a consequence of the Hodge splitting and the Lefschetz fixed-point theorem:

Theorem 3.1. *Let $V(\mathbb{R}) \subset \mathbb{RP}^n$ be an algebraic variety whose complexification $V \subset \mathbb{P}^n$ is an irreducible variety in codimension m . Assume that the singular locus of V has codimension at least $2r + 1$ in V and that, for a generic $L \in \text{Gr}(m + 2r + 1, \mathbb{C}^{n+1})$, the (topological) Euler characteristic $\chi(V \cap L)$ is odd. Then, for any $P \in \text{Gr}(m + 2r + 1, \mathbb{R}^{n+1})$, $V(\mathbb{R}) \cap P \neq \emptyset$.*

In our case, by choosing V as the projectivization of the appropriate degeneracy locus (namely the 5×5 complex matrices with rank 3 or less), and L as a generic 8-dimensional projective space, we show that the Euler characteristic of $V \cap L$ is odd, hence deducing that in the space of 5×5 hermitian matrices with rank not bigger than 3 there are 8 linearly independent elements. This statement is equivalent to say that $d_{5,4} \leq 8$. About the calculation of $\chi(V \cap L)$, the arguments we use are pretty standard ones and essentially concern Schubert calculus of Grassmann spaces and Chern classes of projective bundles.

3.1 Chern classes of determinantal varieties.

Endow the space of complex $q \times q$ matrices $M_q(\mathbb{C})$ with the real structure given by the (antiholomorphic) involution $A \mapsto \overline{A}^t$ (where the bar stands for complex conjugation and t for transposition); with this choice, the real part of $M_q(\mathbb{C})$ is the space $\mathbb{H}_q$ of hermitian matrices. Moreover, the real structure restricts to a real structure over the irreducible affine variety $V_{q,m}^a = \{A \in M_q(\mathbb{C}) \mid \text{rank}(A) \leq m\}$ and its real part $V_{q,m}^a(\mathbb{R})$ is exactly the complement in $\mathbb{H}_q$ of the set $\mathbb{H}_{q,m+1}$ defined in the previous section.

Consider now the projectivizations $\mathbb{P}^{q^2-1} = \mathbb{P}(M_q(\mathbb{C}))$ and $V_{q,m} = \mathbb{P}(V_{q,m}^a)$, and set

$$D_{q,m} = \min\{\dim P \mid V_{q,m}(\mathbb{R}) \cap P \neq \emptyset \text{ for any linear } P \subset \mathbb{RP}^{q^2-1}\}.$$

Proposition 3.1.1. *Let $d_{q,m}$ be as in the previous section. Then $d_{q,m+1} = D_{q,m}$.*

Proof. This should be clear since, by definition of $D_{q,m}$, there is a linear space of projective dimension $D_{q,m} - 1$ entirely included in $\mathbb{P}(\mathbb{H}_{q,m+1}) = \mathbb{R}\mathbb{P}^{q^2-1} - V_{q,m}(\mathbb{R})$. $\square$

We would like, now, to make use of theorem 3.1: we set $V = V_{5,3} \subset \mathbb{P}^{24}$ with codimension 4 and degree 50; its singular locus is $V_{5,2}$ with codimension 5 in $V_{5,3}$ (see e.g. [5]). Our goal is showing that $D_{5,3} \leq 8$, hence the only thing to prove is that, given a generic $L = \mathbb{P}^8 \subset \mathbb{P}^{24}$, one has:

$$\chi(V_{5,3} \cap \mathbb{P}^8) \equiv 1 \pmod{(2)}. \quad (4)$$

To this purpose, consider the following resolution of $V_{5,3}$ with a smooth variety Y :

$$\begin{array}{ccccc} O(1) & \supset & O(1)|_Y & \simeq & \mathcal{O}_{\mathbb{P}}(1) \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{P}(M_5(\mathbb{C})) \times Gr(3, \mathbb{C}^5) & \supset & Y = \{([A], W) \mid \text{Im} A \subset W\} & \simeq & \mathbb{P}(S^{\otimes 5}) \\ & \swarrow \pi_1 & & \searrow \pi_2 & \downarrow \\ & V_{5,3} & & & Gr(3, \mathbb{C}^5). \end{array}$$

Here, $S \rightarrow Gr(3, \mathbb{C}^5)$ is the tautological bundle, $\mathbb{P}(S^5)$ the projective bundle associated to $S^{\otimes 5}$ and $\mathcal{O}_{\mathbb{P}}(1)$ and $O(1)$ are the duals to the tautological bundles over their respective projective bundles. Hyperplane sections of $O(1)|_Y$ are exactly the same of those of $\mathcal{O}_{\mathbb{P}}(1)$. Remark that, outside $\pi_1^{-1}(V_{5,2})$, Y is formed by couples of type $([A], \text{Im} A)$ hence π_1 is generically 1-1; remark also that the isomorphism $Y \xrightarrow{\sim} \mathbb{P}(S^5)$ is given by the map $([A], W) \mapsto A \in \text{Hom}(\mathbb{C}^5, W) \simeq W \otimes \mathbb{C}^{*5} \simeq W^{\otimes 5}$.

For a vector bundle $F \rightarrow B$, let us denote its Chern classes by $c_i(F) \in H^{2i}(B, \mathbb{Z})$ and its Chern polynomial $\sum_{k=0}^{\infty} c_k(F)t^k$ by $c_t(F)$.

Computing $\chi(V_{5,3} \cap \mathbb{P}^8)$ is the same as computing $\chi(Y \cap \bigcap_{i=1}^{16} H_i)$ with H_i generic hyperplanes defined by sections of $O(1)$; since $Z = Y \cap \bigcap_{i=1}^{16} H_i$ is a smooth complex 4-dimensional manifold, denoting by T_Z its tangent bundle and by $[Z] \in H_8(Z, \mathbb{Z})$ its fundamental class, we get

$$\chi(Z) = c_4(T_Z)[Z].$$

Following [14] and [15] we have:

Proposition 3.1.2. *Let $i: Z \rightarrow Y$ be the embedding and $h = c_1(O(1)|_Y)$; then*

$$c_t(T_Y) = i^* c_t(T_Y|_Z)(1 + ht)^{-16}$$

and, denoting by e_4 the coefficient of t^4 in $c_t(T_Y)(1 + ht)^{-16}$:

$$\chi(Z) = h^{16} e_4[Y]. \quad (5)$$

Proof. The first equality is a consequence of the exact bundle sequence over Z : $0 \rightarrow T_Z \rightarrow i^* T_Y|_Z \rightarrow \bigoplus_{i=1}^{16} N_{H_i}|_Z \rightarrow 0$ in which N_{H_i} is the normal bundle of $H_i \subset Y$. The second one follows as in ([15, Ch. 9.2]) since (the restriction of) h represents a submanifold in $Y \cap H_1 \cap \dots \cap H_k$ for every k . $\square$

The diffeomorphism $Y \simeq \mathbb{P}(S^5)$ allows us to compute $c_t(T_Y)$ by means of the exact bundle sequences

$$\begin{aligned} 0 \rightarrow T_{\mathbb{P}(S^5)/G} \rightarrow T_{\mathbb{P}(S^5)} \rightarrow p^*T_G \rightarrow 0 \\ 0 \rightarrow \mathbb{C} \rightarrow S^5 \otimes \mathcal{O}_{\mathbb{P}}(1) \rightarrow Q \otimes \mathcal{O}_{\mathbb{P}}(1) = T_{\mathbb{P}(S^5)/G} \rightarrow 0 \end{aligned}$$

where the first one comes from the projection $p : \mathbb{P}(S^5) \rightarrow G = G(3, \mathbb{C}^5)$ and the second one is the tautological sequence over $\mathbb{P}(S^5)$ tensorised with $\mathcal{O}_{\mathbb{P}}(1)$. Working out calculations we find (see [14]):

$$c_t(T_{\mathbb{P}(S^5)}) = c_t(T_G)c_t(T_{\mathbb{P}(S^5)/G}) = c_t(T_G) \left(\sum_{j=0}^{15} c_j(S^5)t^j(1+ht)^{15-j} \right). \quad (6)$$

3.2 Computation of the Euler characteristic (modulo 2).

We now dispose of all the needed tools to complete the proof of the statement 4. Remark that universal coefficient theorem implies that $H^*(B, \mathbb{Z}) \otimes \mathbb{Z}_2 = H^*(B, \mathbb{Z}_2)$, where B is either $G(3, \mathbb{C}^5)$ or $\mathbb{P}(S^5)$. Thanks to this, since we are only interested to the parity of our objects, we will perform any computation in $H^*(\cdot, \mathbb{Z}_2)$; moreover all polynomials in the Chern classes will be 1.1 truncated according to our real necessities.

Let us denote by c_i the Chern classes of S .

Proposition 3.2.1. *Modulo 2, we have:*

1. $c_t(S^5) = 1 + c_1t + c_2t^2 + c_3t^3 + c_1^4t^4 + c_1^5t^5 + c_1^4c_2t^6$;
2. $c_t(T_G) = 1 + c_1t + c_2t^2 + c_3t^3 + c_2^2t^4 + c_1c_2^2t^5 + (c_1^4c_2 + c_3^2)t^6$;
3. the ring $H^*(\mathbb{P}(S^5), \mathbb{Z}_2)$ is $\mathbb{Z}_2[c_1, c_2, h]$ together with the relations

$$h^{15} = \sum_{j=1}^{15} c_j(S^5)h^{15-j} \quad c_1c_2^2 = 0 \quad c_1^4 + c_1^2c_2 + c_2^2 = 0 \quad (7)$$

Proof. 1. Follows directly from $c_t(S^5) = c_t(S)^5$.

2. Follows from the sequence $0 \rightarrow S \otimes \tilde{S} \rightarrow \tilde{S}^5 \rightarrow Q \otimes \tilde{S} = T_G \rightarrow 0$ where $\tilde{S}$ is the dual of S and $c_t(\tilde{S}) = c_{-t}(S) = c_t(S)$ since we are working in $\mathbb{Z}_2$. Hence: $c_t(T_G) = c_t(S \otimes \tilde{S})c_t(S)^5$.

3. (For a more detailed treatment, see[8]).

The structure of the cohomology ring of a projective bundle $\mathbb{P}(E) \rightarrow B$ is well known: $H^*(\mathbb{P}(E)) \simeq H^*(B)[h]/(h^r + \sum_1^r c_i(E)h^{r-i})$ with h the first Chern class of the canonical bundle over $\mathbb{P}(E)$ and $r = \text{rank}(E)$.

We have that $H^*(Gr(k, \mathbb{C}^n), \mathbb{Z}_2) = \mathbb{Z}_2[c_1, \dots, c_k]/(s_{n-k+1}, \dots, s_n)$ where c_i are the Chern classes of the tautological bundle $S \rightarrow Gr(k, \mathbb{C}^n)$ and s_i those of the quotient bundle $\mathbb{C}^n/S$.

In our context $s_3 = c_1^3 + c_3$, $s_4 = c_1^4 + c_1^2c_2 + c_2^2$ and $s_5 = c_1^5 + c_1^2c_3 + c_1c_2^2$; the statement follows from the elimination of c_3 and simplification of the expression of s_5 .

□

Let e_4 be the coefficient of t^4 in $c_t(T_{\mathbb{P}(S^5)})(1+ht)^{-16}$;

Lemma 3.2.2. $e_4 = h^4 + c_1 h^3 + c_1^2 h^2 + c_2 h^2 + c_1^3 h + c_1 c_2 h + c_1^4$ modulo 2.

Proof. From equation (6) we have

$$c_t(T_{\mathbb{P}(S^5)})(1+ht)^{-16} = c_t(T_G)(1+ht)^{-1} \left(\sum_{j=0}^{15} c_j(S^5) \left(\frac{t}{1+ht} \right)^j \right);$$

applying 3.2.1 and truncating polynomials to the 4th degree, this becomes

$$(1+c_1 t+c_2 t^2+c_1^3 t^3+c_2^2 t^4)(1+ht+h^2 t^2+h^3 t^3+h^4 t^4) \sum_{j=0}^4 c_j(S^5)(t+ht^2+h^2 t^3+h^3 t^4)^j.$$

The statement follows simply by working out calculations and by deleting couples of equal terms. $\square$

Proposition 3.2.3. *The class $h^{16}e_4$ generates the top cohomology of $H^*(\mathbb{P}(S^5), \mathbb{Z}_2)$.*

Proof. The proof of this statement is almost entirely based on the third point of 3.2.1. The second and third relation of (7) imply that $H^{12}(Gr(3, \mathbb{C}^5), \mathbb{Z}_2)$ is generated by any degree 6 monomial in c_1, c_2 , but $c_1^2 c_2^2$.

We claim that $H^{40}(\mathbb{P}(S^5), \mathbb{Z}_2)$ is generated by gh^{14} where g is any generator of $H^{12}(Gr(3, \mathbb{C}^5), \mathbb{Z}_2)$. This can be seen directly by reducing the degree of h in any monomial of type $m(c_1, c_2)h^{14+k}$ with the aid of the first relation of (7). As an example, consider $c_1^2 c_2 h^{16}$:

$$\begin{aligned} c_1^2 c_2 h^{16} &= c_1^2 c_2 h^{15} = c_1^2 c_2 h (c_1 h^{14} + c_2 h^{13} + \text{higher order terms in } c_1, c_2) = \\ &= c_1^3 c_2 h^{15} + c_1^2 c_2^2 h^{14} = c_1^3 c_2 (c_1 h^{14} + \dots) + 0 = c_1^4 c_2 h^{14}. \end{aligned}$$

Granting this, one finds (by analogous calculations) that the first and the three last terms in the expression of $h^{16}e_4$ given by 3.2.2 vanish, while the three others do not. Hence, $h^{16}e_4$ generates $H^{40}(\mathbb{P}(S^5), \mathbb{Z}_2)$. $\square$

Corollary 3.2.4. *If L is a generic 8-dimensional linear subspace of $\mathbb{P}^{24}$, the euler characteristic $\chi(L \cap V_{5,3})$ is odd.*

Proof. By equation (5):

$$\chi(L \cap V_{5,3}) = \chi(Z) = h^{16}e_4[\mathbb{P}(S^5)]$$

and the latter term is not zero modulo 2 thanks to 3.2.3 and Poincaré duality between $H^{40}(\mathbb{P}(S^5), \mathbb{Z})$ and $H_{40}(\mathbb{P}(S^5), \mathbb{Z})$. $\square$

Remark 3.2.5. *The following matrix, depending on 7 real parameters, shows that $d_{5,4} = D_{5,3} \geq 7$, since $\text{rank} A \geq 4$ unless $\alpha = z = u = w = 0$.*

$$A(\alpha, z, u, w) = \begin{pmatrix} \alpha & z & u & w & 0 \\ \bar{z} & \alpha & \bar{w} & -\bar{u} & 0 \\ \bar{u} & w & -\alpha & z & 0 \\ \bar{w} & -u & \bar{z} & -\alpha & z \\ 0 & 0 & 0 & \bar{z} & 0 \end{pmatrix} \quad \alpha \in \mathbb{R}, \quad z, u, w \in \mathbb{C}.$$

To see this, let A_k be the submatrix obtained from A by elimination of the k -th row and column; A_3 is invertible unless $z = 0$ or $|\alpha| = |z|$. When $z = 0$, $A_5^2 = (\alpha^2 + |u|^2 + |w|^2)I$ (see also [2]); when $|\alpha| = |z|$, $\det A_1 = |z|^2(|z|^2 + |w|^2)$.

References

- [1] J. Adams. *Vector fields on spheres*. Ann. of Math. (2) **75** (1962) 603-632.
- [2] J. Adams, P. Lax, and R. Phillips. *On matrices whose real linear combinations are non-singular*. Proc. Amer. Math. Soc. **16** (1965) 318-322.
- [3] J. Amòros and I. Bauer. *On the number of defining relations for nonfibered Kähler groups*. Internat. J. Math. **11** (2000), 285-290 .
- [4] J. Amòros, M. Burger, K. Corlette, K. Kotschick and D. Toledo. *Fundamental groups of compact Kähler manifolds*. Mathematical Surveys and Monographs, **44** (1996) A.M.S., Providence, RI.
- [5] E. Arbarello, M. Cornalba, P. Griffiths, J. Harris. *Geometry of algebraic curves*. Grundlehren der Mat. Wissenschaften (vol. 267), Springer, 1985.
- [6] M Barja, C. Naranjo and G. Pirola. *On the topological index of irregular varieties*. preprint.
- [7] W. Barth, C. Peters and A. Van de Ven. *Compact Complex Surfaces*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer Verlag, 1984.
- [8] R. Bott, W. Tu. *Differential forms in algebraic topology*. Graduate text (vol. 82), Springer, 1982
- [9] F. Campana. *Remarques sur les groupes de Kähler nilpotents*. Ann.Sci Ecole Norm. Sup. (4) **28** (1995), no. 3, 307-316.
- [10] F. Catanese. *Moduli and classification of irregular Kähler manifolds (and algebraic varieties) with Albanese general type fibrations*. Invent. Math. **104** (1991), 263-289.
- [11] O. Debarre. *Théorèmes de connexité et variétés abéliennes*. Amer. J. Math. **117** (1995), no 3, 787-805.
- [12] P. Deligne, P. Griffiths, J. Morgan and D. Sullivan. *Real homotopy theory of Kähler manifolds*. Invent. Math, **29** (1975) 245-274.
- [13] D. Falikman, S. Friedland and R. Loewy. *On spaces of matrices containing a nonzero matrix of bounded rank*. Pacific J. Math., **207** (2002) no. 1, 157-176.
- [14] S. Friedland and A. Libgober. *Generalizations of the odd degree theorem and applications*. Israel J. Math. **136** (2003), 353-371.
- [15] F. Hirzebruch. *Topological methods in algebraic geometry*. Grundlehren der mat., Wissenschaften (vol. 131), Springer, 1966.

ANDREA CAUSIN

Dipartimento di Matematica, Università “La Sapienza” di Roma
P.le Aldo Moro 2, 00185 Roma, Italia
causin@mat.uniroma1.it

GIAN PIETRO PIROLA

Dipartimento di Matematica, Università di Pavia
via Ferrata 1, 27100 Pavia, Italia
pirola@dimat.unipv.it